

**A supplementary note on *Lathys* (*Scotolathys*)
punctosparsa Oi**

Ryoji Oi

Faculty of the Science of Living, Osaka City University

In my previous note (Acta Arachnologica Vol. 14, No. 2, p. 45, 1957) *Lathys punctosparsa* was described on the female specimen from Buttsu-ji, Hiroshima Prefecture. As two males and three females were captured by myself at Yata, Osaka City on October 10, 1959, in the present paper I give a supplementary description on the male specimen.

The structure and colour of the male are similar to those of the female, but sometimes there are several rather large faintly dark markings on the dorsum and sides of the abdomen instead of the many procurved slender stripes consisting of small black spots on the abdomen as mentioned in my previous note.

Palpal structures of the male are as showing in figures.

The measurements are as follows. (m.m.)

♂ : Body :	Length			Width	
	Total	Cephalothorax	Abdomen	Cephalothorax	Abdomen
	2.13	1.00	1.13	0.75	0.82

♂ : Legs :	Femur	Patella & Tibia	Metatarsus	Tarsus	Total
I	1.25	1.38	0.93	0.62	4.18
II	1.00	1.18	0.83	0.54	3.55
III	0.94	1.00	0.81	0.50	3.25
VI	1.19	1.28	0.94	0.63	4.04




Fig. 1 Ventromedial aspect of left palp.

Fig. 2 Latetel aspect of left palp.